

THE SILURIAN BRACHIOPOD *STEGERHYNCHUS*

by BRIAN JONES

ABSTRACT. The brachiopod *Stegerhynchus* Foerste, 1909 occurs in Silurian strata in Canada, the U.S.A., the U.S.S.R., Gotland, and the United Kingdom. Review of the literature shows that *Rhynchonella whitii praecursor* Hall, 1863 is the true type species of *Stegerhynchus*. Although *Stegerhynchus* and *Ferganella* Nikiforova, 1937 are morphologically very similar, they can be separated because *Stegerhynchus* has an open notothyrial cavity that houses an elongate cardinal process based on the notothyrial platform, whereas *Ferganella* has a septal process extending anteriorly from the posteriorly conjunct hinge plates. The septal process joins the cardinal process distally. *Stegerhynchus* seems to be most common in Wenlock and Ludlow strata and less common in Llandovery and Pridoli strata. *Ferganella* appears to be of Pridoli/Lower Devonian age. Species such as *S. borealis* and *S. diodonta* are relatively long-ranging while other species such as *S. angaciensis* have a relatively short range. *S. borealis* is relatively common in the Upper Silurian Read Bay Formation of the Canadian Arctic islands, whereas *S. angaciensis* is relatively rare, being found only at the base of member C of the Read Bay Formation on the east coast of Cornwallis Island.

DURING a detailed study of the Ludlovian and Pridolian brachiopods from the Read Bay Formation of Arctic Canada it became apparent that there was considerable confusion surrounding the definition of *Stegerhynchus* and *Ferganella*. The literature covering these genera contains many conflicting statements. Amsden (1974, p. 67) commented that '... the distinction between *Stegerhynchus* and *Ferganella* is not at all clear ...'. This study outlines some of the main problems surrounding the genus *Stegerhynchus*. It is based on a review of all the North American species assigned to *Stegerhynchus* or *Ferganella* with a detailed analysis of the literature. Reference is also made to the European material for comparative purposes. In addition the rhynchonellids from the Read Bay Formation of Arctic Canada are described. I am particularly indebted to Dr. M. G. Bassett of the National Museum of Wales who provided me with information pertaining to European specimens of *Stegerhynchus*. Dr. Bassett also allowed me to see type material of *Ferganella turkestanica*, *F. borealis*, *S. decemplicatus angaciensis*, and *F. diodonta* which he had spent so much time and effort to collect and borrow. Dr. Bassett is currently studying the *Stegerhynchus* and *Ferganella* from Europe and is further investigating the taxonomic relationships in this group of brachiopods.

In this paper the following abbreviations are used for the numbering of type material: GSC = Geological Survey of Canada repository, Ottawa; UCLA = University of California, Los Angeles; OU = University of Oklahoma; UI-RX = University of Illinois, Rowley collection; NMW = National Museum of Wales; Br = Naturhistoriska Riksmuseet, Stockholm, and ETA = Eesti Teadusti Akademy; UA = University of Alberta.

TAXONOMIC HISTORY OF *STEGERHYNCHUS*

Foerste (1909, p. 98) introduced the name *Stegerhynchus* 'to distinguish the species typified by *Rhynchotreta whitii praecursor*, from the more typical species of *Rhynchotreta* ...'. Since Foerste (1909) did not designate a type species Schuchert and LeVene (1929) selected *Rhynchonella whitii* as the genoelectotype of *Stegerhynchus*. Kozłowski (1929, p. 146) listed *Stegerhynchus* as a synonym of *Stenocisma* Conrad (1839), reasoning that the external and internal structures of *Stegerhynchus* as described by Foerste (1909) were exactly the same as those described for *Stenocisma* by Conrad (1839) (Kozłowski 1929, p. 148). However, Ager (1965, p. H629) included *Stenocisma* Conrad (1839) (not *Stenocisma* Hall, 1847; *Stenocisma* Hall and Clarke, 1894 or *Stenocisma* Grabau and Shimer 1907) in the family Stenocismatidae Oehlert (1883), since the type species, *Terebratulata schlotheimi*,

Von Buch (1835), is a rhynchonelliform rhynchonellid with a well-developed stolidium. *Stegerhynchus* cannot be considered a synonym of *Stenochisma* Conrad (1839).

Shimer and Shrock (1944, p. 309), ignored the work of Schuchert and LeVene (1929) and listed *R. whitii praecursor* as the type species of *Stegerhynchus* Foerste 1909. Cooper (1955, p. 54) and Cocks (1978, p. 147) concluded that *R. whitii* Hall was the type species of *Stegerhynchus* since Schuchert and LeVene (1929, p. 42) had designated it as such. The *Treatise* lists *R. whitii praecursor* (= *S. praecursor*) as the type species of *Stegerhynchus*. The choice between the two species is critical because it has been shown by Cooper (1955, p. 54) and Amsden (in Amsden and Boucot, 1958, pp. 154-155) that *R. whitii praecursor* has a cardinal process while *R. whitii* does not. According to the *Treatise*, rhynchonellids in the subfamily Rhynchotrematinae have a cardinal process while those in the subfamily Trigonirhynchiidae do not. If *R. whitii praecursor* is the type species, *Stegerhynchus* would be assigned to the subfamily Rhynchotrematidae. If *R. whitii* is the type species, *Stegerhynchus* would be placed in the subfamily Trigonirhynchiidae. The problem of the type species of *Stegerhynchus* centres around the genolecotype chosen by Schuchert and LeVene (1929). Foerste's (1909) paper suggests that their choice was incorrect for the following reasons:

(a) Foerste (1909, p. 98) stated: 'To distinguish the species typified by *Rhynchotretra whitii praecursor*, from the more typical species of *Rhynchotretra*, possessing an acuminate beak, long broad flattened sides, and a median depression along the posterior parts of the brachial valve, the term *Stegerhynchus* may be employed.' It must be assumed that *Rhynchotretra* was mistakenly used for *Rhynchonella* since it is clear from previous references to *whitii praecursor* (Foerste 1909, p. 96) that it belonged to *Rhynchonella*, not *Rhynchotretra*.

(b) *R. whitii praecursor* was fully described by Foerste (1909, pp. 96-97) while *R. whitii* was only quoted in a list of species considered congeneric with *R. neglecta* and *R. indianensis*.

(c) *R. whitii praecursor* and *R. neglecta cliffonensis* were both figured by Foerste (1909, pl. III, figs. 47A, B, C and 48A, B, C respectively) while *R. whitii* was not figured.


(d) *R. whitii praecursor* is the first described form linked with the name *Stegerhynchus*.

Cooper (1955, p. 54) considered that Schuchert and LeVene (1929) had chosen *R. whitii* as the type species since it was the first name in the list of species considered congeneric with the new varieties named by Foerste. From Foerste's (1909, p. 98) concluding statement it is apparent that he considered *R. whitii praecursor* typical of *Stegerhynchus*. Also, *R. whitii praecursor* was the first name linked with *Stegerhynchus* and was better described and illustrated than *R. whitii* and thus should be given preference.

In view of the above considerations and the opinion given by Amsden (1968, p. 62) it is clear that *R. whitii praecursor* should be taken as the type species of *Stegerhynchus*.

COMPARISON WITH CLOSELY RELATED GENERA

Nikiforova (1937, pp. 77-78) erected the genus *Ferganella*, based on specimens from the northern slope of the Altai Range (Fergara). *F. turkestanica*, the type species, has a moderate to large, subrounded to subtriangular shell of variable convexity with a distinct fold and sinus. In the pedicle valve it has massive teeth supported by dental plates. In the brachial valve the massive, thick septum supports the notothyrial cavity in which the elongate cardinal process is housed. Joining the posteriorly conjunct hinge plates is a septal process which extends anteriorly before joining the cardinal process in a distal position (text-fig. 1). The crura extend from the inner edges of the hinge



TEXT-FIG. 1. Serial sections showing internal structures of *Ferganella turkestanica*. Diagrams based on illustrations in Nikiforova (1937, pl. 7, figs. 15 and 16). Exact positions relative from posterior not known, but *a* is closer than *b*.

plates. Nikiforova (1937, p. 37) transferred *R. borealis* Schlotheim to *Ferganella* because of the great similarity of the internal structures to those found in *F. turkestanica*.

Nikiforova (1937, p. 37) did not compare *Ferganella* directly with *Stegerhynchus*. Rather she compared *Stenochisma* (considered the senior synonym of *Stegerhynchus* by Kozłowski 1929) with *Ferganella*, noting that *Stenochisma* lacked a median septum which is well developed on *Ferganella*. Although *Ferganella* differs from *Stenochisma* according to the criteria listed by Nikiforova (1937, p. 78) it does not follow that the same differences serve to separate *Stegerhynchus* from *Ferganella* as *Stegerhynchus* is not a synonym of *Stenochisma*. Externally, two specimens of *F. turkestanica* from a locality on the River Djalkin in Fergana (NMW 76.9G.56 and NMW 76.9G.57; shown to me by Dr. M. G. Bassett) are very similar to many of the species herein assigned to *Stegerhynchus*. However, comparison of the internal structures of *Ferganella* and *Stegerhynchus* shows that *Stegerhynchus* has an open notothyrial cavity, posteriorly disjunct hinge plates, and no septal process, whereas *Ferganella* has posteriorly conjunct hinge plates and a septal process which joins the cardinal process anteriorly. These internal structures of *Ferganella* are evident from the original description of *F. turkestanica* (Nikiforova 1937), from the illustrations of the internal structures of *F. turkestanica* (text-fig. 1) and from an original section of *F. turkestanica* shown to me by Dr. M. G. Bassett. These differences separate the two genera.

Rzhonsnitskaya (1959, p. 27) defined *Stegerhynchella* using *S. decemplicatus angaciensis* (Chernyshev) as the type species. However, Schmidt and McLaren (1965, p. H556) and Lenz (1970, p. 488) suggested that *Stegerhynchella* is probably a synonym of *Stegerhynchus*. Inspection of twenty-two specimens of *S. decemplicatus angaciensis* (Chernyshev) from the topotype locality in the Elegest Formation on the River Elegest, central Tuva, U.S.S.R., confirms this. Amsden recently named the new genus *Stegerhynchops* for non-lamellose rhynchotrematids that lacked dental plates. The absence of dental plates clearly separate it from *Ferganella* and *Stegerhynchus*.

DISCUSSION


Until now *Stegerhynchus* and *Ferganella* have been maintained as separate genera even though the criteria for separating the two have not been clearly delineated. Since Cooper (1955, p. 54) considered *R. whitii* the type species of *Stegerhynchus*, he assigned *R. (Stegerhynchus) whitii praecursor* Foerste, 1909 and *R. (Stegerhynchus) neglecta cliftonensis* Foerste, 1909 to *Ferganella* because they both possessed a cardinal process. Havlicek (1961, pp. 80-91) and Kulkov (1963, 1967) followed Cooper's example, taking *R. whitii* as the type species of *Stegerhynchus*, and therefore considered that *Stegerhynchus* did not have a cardinal process. Johnson and Reso (1964, p. 80) assigned rhynchonellids from the Sevy Dolomite of Nevada to *Ferganella*, on the basis that their external form was closer to *Ferganella* than *Stegerhynchus*. Lenz (1970, p. 488) assigned rhynchonellids from Prongs Creek to *Ferganella* rather than *Stegerhynchus* because of their external characteristics and because of the '... presence of the median septum, which though variable in height and development, is normally well developed'. Although there is the implication that *Ferganella* and *Stegerhynchus* can be separated by virtue of the size and form of the median septum, the differences were not specified. Study of the various species of *Stegerhynchus* and *Ferganella* revealed no readily discernible differences at the generic level (Table 1).

The concept that the nature of the median septum could be used for separating the two genera was expanded by Johnson, Boucot, and Murphy (1976, p. 64) who considered that *Rhynchotrema* and *Ferganella* both have a true median septa in their brachial valve while *Stegerhynchus* does not. This statement was made without supportive evidence or illustrations, but Johnson (1977, written comm.) considers that *Stegerhynchus* has a myophragm rather than a median septum. Although Johnson *et al.* (1976) considered the dichotomy between the genera to be quite rigid, study of the pertinent descriptions, illustrations, and, where possible, type material has shown that the length, height, and form of the median elevation is highly variable (Table 1). In *Stegerhynchus* the length of the median elevation ranges from 43 to 65% of valve length compared to 36 to 63% of valve length in *Ferganella* (Table 1). There is no difference in the length of the median elevations of species assigned to

TABLE 1. Summary of main morphological attributes of the type specimens of *Stegerhynchus angaciensis*, *S. claritense*, *S. concinna*, *S. antiqua*, *F. cf. F. lincolnensis*, *F. chattertoni*, *F. didonta*, *F. borealis*, and *F. turkestanica*. Abbreviations used are: C = complete; BV = Brachial valve; PV = Pedicle valve; PAS = Posterior articulated shell.

Species Reference	Type Number	Valves present	Number of ribs in sulcus	Number of ribs on fold	Number of ribs on flanks of PV		Type Number	Valves present	Number of ribs in sulcus	Number of ribs on fold	Number of ribs on flanks of PV	
					P	M					P	M
Species Reference	STEGEHRYNCHUS ANGACIENSIS LENZ, 1970			Chernyshev		FERGANELLA cf. F. LINCOLNENSIS Johnson LENZ, 1970						
	GSC 25007	C	1	2	2	1	GSC 25000	C	3	4	3	1
	GSC 25008	C	1	2	3	2	GSC 25001	BV	-	4	3	1
	GSC 25009	C	1	2	4	1	GSC 25002	C	3	4	3	2
	GSC 25010	BV	-	2	-	-	GSC 25003	PAS	-	-	-	-
	GSC 25011	PAS	-	2	3	1	GSC 25004	PV	2	-	3	1
	GSC 25012	PV	1	-	4	1	GSC 25005	BV	-	4	-	-
GSC 25013	BV	-	-	-	-	GSC 25006	BV	-	-	-	-	
Species Reference	STEGEHRYNCHUS ? sp. LENZ, 1977					FERGANELLA cf. F. LINCOLNENSIS Johnson and Reso, 1964						
	GSC 48075	C	1	2	3	1	UCLA 35054 (HY)	C	2	3	4	0
	GSC 48076	BV	-	3	-	-	UCLA 35050 (HOL)	C	3	4	4	1
GSC 48066	C	2	3	3	1	UCLA 35053 (P)	C					
Species Reference	STEGEHRYNCHUS CLARITENSE AMSDEN, 1968			Amsden		FERGANELLA CHATTERTONI LENZ, 1977						
	OU 6384 (HOL)	C	1	2	4	1	GSC 48067	C	3	4	3	1
OU 6385	C	1	2	2	1	GSC 48068	C	3	4	3	1	
Species Reference	STEGEHRYNCHUS CONCINNA AMSDEN, 1974			(Savage)		FERGANELLA DIDONTA (Gelman) BASSETT AND COCKS, 1974						
	OU 6762	C	3	4	4	1	GSC 48069	-	-	-	-	
	OU 6752	C					GSC 48070	PV	3	-	3	1
	RX-855 (LECT)	C	2	3	5	-	GSC 48071	PV	3	-	3	1
	RX-306	PVI	2	-	4	1	GSC 48072	BV	-	-	-	
	RX-307	BVI	-	3	4	1						
	RX-308	C	2	3	6	4	RMS Br 128B (LECT)	C	1	2	4	1
					4	4						
	RX-309	C	2	3	4	1						
	RX-310	BVI	-	-	-	-						
	RX-311	C	3	4	5	1						
	RX-312	C	1	2	3	1						
	RX-313	C	3	4	5	1						
	RX-314	PVI	3	-	4	1						
	RX-315	C	2 (?)	3 (?)	5	1						
	RX-316	C	2	3	4	1						
	RX-317	C	3	4	3	1						
RX-318	C	2	3	5	1							
RX-319	C	2	3	5	1							
Species Reference	STEGEHRYNCHUS ? ANTIQVA AMSDEN, 1974			(Savage)		FERGANELLA TURKESTANICA Nikiforova NIKIFOROVA, 1937						
	RX-4784	C	3	4	4	1	P1. VII fig. 10	C	5	6	4	1
	RX-B72a	C	3	4	4	1	P1. VII fig. 11	C	5	6	5	1
	b	C	3	4	3	2	P1. VII fig. 12	C	5 (?)	6	4	(?)
	c	C	4	5	3	1	P1. VII fig. 13	C				Not clear from photograph
Species Reference	STEGEHRYNCHUS ? sp. AMSDEN, 1974			Very poor specimen		FERGANELLA TURKESTANICA var NYMPHAEFORMIS Nikiforova NIKIFOROVA, 1937						
	RX-320	C					P1. VII fig. 17	C	5	6	4	1
	RX-321	C	1	2			P1. VII fig. 18	C	7	8	6	1

Stegerhynchus or *Ferganella*. Similarly, the height and development of the structure is also highly variable. For example, in *S. angaciensis* described by Lenz (1970) GSC types 25010 and 25013 both have well-preserved median septa. These two specimens show a marked contrast, GSC 25010 (text-fig. 2c) has a relatively short but high median elevation whereas GSC 25013 (text-fig. 2h) has a relatively long but low median elevation. This variability is emphasized further because the relatively high median elevation of GSC 25010 is skewed off on to the right-hand flank of the shell (text-fig. 2e).



TEXT-FIG. 2. Internal structures of *Stegerhynchus*: a, brachial valve interior of *S. concinna* showing open notothyrial cavity housing elongate cardinal process. Medium septum supports notothyrial cavity. Specimen UI-RX-307; b, brachial valve interior of *S. concinna* showing same structures as in fig. 2a. Note medium septum is more prominent. Specimen UI-RX-310; c, pedicle valve interior of *S. concinna* showing dental plates and teeth. Specimen UI-RX-306; d, posterior portion of articulated specimen of *F. chattertoni* showing elongate cardinal process in open notothyrial cavity. Note crus extending from inner edges of hinge plates, well-developed medium septum and well-developed teeth in sockets. Specimen GSC 48070; e, brachial valve interior of *S. angaciensis* from Prongs Creek showing 'right-skewed' medium septum and associated asymmetry of right flank of shell. Note open notothyrial cavity. Specimen GSC 25010; f, brachial valve interior of *F. chattertoni* showing open notothyrial cavity, elongate cardinal process, relatively broad medium septum, and curved hinge plates. Specimen GSC 48069; g, brachial valve interior of *F. cf. F. lincolnensis* from Prongs Creek showing open notothyrial cavity, elongate cardinal process, sockets, triangular hinge plates, and relatively broad medium septum. Specimen GSC 25006; h, brachial valve interior of *S. angaciensis* from Prongs Creek. Note difference in size, form, and location of medium septum compared to specimen in fig. 2e. Specimen GSC 25013.

F. lincolnensis described by Lenz (1970) has a median elevation that extends 57 to 63% of shell length. In specimens GSC 25001 and 25005 the median elevation is relatively low and merges imperceptibly with the central inter-rib space of the fold. By contrast, *F. chattertoni*, as defined by Lenz (1977), has a median elevation 36 to 40% valve length. Specimen GSC 48069 (text-fig. 2f) has a relatively high elevation whereas specimens GSC 48072 and 48073 have relatively low median elevations. In specimens GSC 48069 (text-fig. 2f) and GSC 48072 the anterior termination of the median elevation is relatively well defined, whereas in specimen GSC 48073 the termination is imperceptible and difficult to define. Similar variation can also be seen in *S. concinna* and Amsden (1974, p. 66) describes the presence of a median septum in *S. concinna*. Again, there is the implication that the size and form of the median elevation is highly variable and therefore difficult to use as a criterion for separating genera. If such a high degree of variation is evident from a limited number of specimens from widely scattered areas, then it is difficult to argue that a larger sample size would clarify the dichotomy suggested by Johnson *et al.* (1976). One important point to consider is demonstrated by study of specimen GSC 48076 (*Stegerhynchus*?) and an unnumbered specimen associated with specimen

GSC 48076. The brachial valve of the unnumbered specimen is more convex than the brachial valve of specimen GSC 48076 and has a correspondingly higher median elevation; implying that the height of the median septum is controlled by the distance between the floor of the brachial valve and the base of the notothyrial cavity. Therefore, the more convex valves will have the highest median elevation. This factor is probably partly responsible for the high septum in specimen GSC 25010 as opposed to the relatively low septum in specimen GSC 25013.

Summary. Review of the literature has shown that there is considerable confusion surrounding the genus *Stegerhynchus*. Moreover, the distinction between *Stegerhynchus* and *Ferganella* has been particularly difficult to delineate. There have been four main approaches to the latter problem; each with its associated problems, namely:

(a) The presence or absence of a cardinal process: this resulted from confusion over the type species of *Stegerhynchus*. However, since it is now clear that *S. praecursor* is the true type species of *Stegerhynchus*, this criterion cannot be used.

(b) Shell shape: although it has been stated that the external forms of the two genera differ, the differences have not been described in detail. Study of numerous specimens of *Stegerhynchus* suggests that there is a great range of shell shapes within the genus and that separation from another genus such as *Ferganella* on this basis of this criterion would probably prove very unsatisfactory.

(c) Form of median elevation in brachial valve: it has been stated that *Ferganella* has a true median septum whereas *Stegerhynchus* only has a myophragm. Study of all the available material suggests that the form of the median elevation is variable even amongst shells from the same locality. In this respect there does not appear to be a distinct dichotomy between the two genera.

(d) Internal structures: *Stegerhynchus* has an open notothyrial cavity whereas *Ferganella* has posteriorly conjunct hinge plates which partly cover the cavity and a septal process which joins the cardinal process in a distal position. This seems to be the only reliable way of separating *Stegerhynchus* from *Ferganella*.

Bassett (1978, written comm.) has also pointed out that Nikiforova's original specimens of *F. turkestanica* came from strata of Devonian age. Thus, it would appear that *Ferganella* is younger than *Stegerhynchus* which occurs mainly in Silurian strata. In view of the foregoing discussion the species herein considered part of *Stegerhynchus* are assigned on the basis of the last of the internal structures listed above.

SYSTEMATIC PALAEOLOGY

Order RHYNCHONELLIDA Kuhn, 1949

Superfamily RHYNCHONELLACEA Gray, 1848

Family RHYNCHOTREMATIDAE Schuchert, 1913

Subfamily RHYNCHOTREMATINAE Schuchert, 1913

Genus STEGERHYNCHUS Foerste, 1909

- 1909 *Rhynchonella* (*Stegerhynchus*) (*pars*) Foerste, pp. 96-97, pl. III, fig. 47A, B, and C.
 1937 *Stegerhynchus* Foerste, 1909; Chernyshev, p. 29, p. 71, pl. 1, figs. 15-18; text-figs. 1-2.
 1937 *Camarotochia* Hall and Clarke, 1894; St. Joseph, pp. 33-48, figs. 1-5.
 1955 *Stegerhynchus* Foerste, 1909; Borisyak, pp. 46-47, pl. VI, figs. 6-11.
 1955 *Ferganella* Nikiforova, 1937; Cooper, p. 55.
 1959 *Stegerhynchella* Rzhonsnitskaya, p. 27.
 1960 *Stegerhynchella* Rzhonsnitskaya, 1959; Rzhonsnitskaya, Likharev, and Makridin, pl. 43, fig. 7a-d; text-figs. 246-247.
 1960 *Stegerhynchella* Rzhonsnitskaya, 1959; Khalina, p. 102, pl. S-28, fig. 5a-c.
 1960 *Stegerhynchus* Foerste, 1909; Zinchenko and Kulkov, p. 103, pl. S-28, fig. 7.
 1929 *Stenochisma* Conrad, 1839; Kozłowski (*pars*), pp. 146-150.
 1964 *Ferganella* Nikiforova, 1937; Johnson and Reso, p. 80, pl. 19, figs. 5-12.
 1967 *Ferganella* Nikiforova, 1937; Kulkov, pp. 75-76, text-fig. 31; pl. XII, figs. 7-11.

- 1968 *Stegerhynchus* Foerste, 1909; Amsden, pp. 61–63, text-fig. 47; pl. 17, fig. 1a–k.
 1970 *Stegerhynchus* Foerste, 1909; Lenz, pp. 488–489, pl. 85, figs. 16–29.
 1970 *Ferganella* Nikiforova, 1937; Lenz, pp. 487–488, pl. 85, figs. 1–15.
 1974 *Ferganella* Nikiforova, 1937; Bassett and Cocks, p. 26, pl. 8, fig. 2A–D.
 1974 *Stegerhynchus* Foerste, 1909; Amsden, pp. 66–68, text-figs. 41–42; pl. 14, figs. 3–4; pl. 15, figs. 1–4.
 1976 *Ferganella* Nikiforova, 1937; Sheehan, p. 729, pl. 5, figs. 1–5.
 1976 *Ferganella* Nikiforova, 1937; Smith, p. 29, text-fig. 19; pl. 7, figs. 20–25.
 1976 *Stegerhynchus* Foerste, 1909; Johnson *et al.*, pp. 64–65, pl. 47, figs. 1–12.
 1977 *Ferganella* Nikiforova, 1937; Lenz, p. 1542, pl. 8, figs. 1–16.
 1977 *Ferganella* Nikiforova, 1937; Lenz, pp. 87–88, pl. 17, figs. 1–6 and 11.
 1977 *Stegerhynchus* Foerste, 1909; Lenz, p. 88, pl. 17, figs. 7–10, 12–29.
 1978 *Stegerhynchus* Foerste, 1909; Amsden, pl. 11, figs. 1–9.
 non 1955 *Stegerhynchus* Foerste, 1909; Cooper, pp. 54–55.
 non 1961 *Stegerhynchus* Foerste, 1909; Havlicek, p. 80.
 non 1963 *Stegerhynchus* Foerste, 1909; Kulkov, pp. 45–46.
 non 1964 *Stegerhynchus* Foerste, 1909; Drot, p. 103, pl. 19, figs. 6–8.
 non 1967 *Stegerhynchus* Foerste, 1909; Gratsianova, p. 68, pl. 6, fig. 6.
 non 1970 *Stegerhynchus* Foerste, 1909; Alekseeva, Gratsianova, Elkin, and Kulkov, p. 70, pl. 7, figs. 10–11.

Type species. *S. praecursor* Foerste, 1909 (= *Rhynchonella* (*Stegerhynchus*) *whitii praecursor* of Foerste, 1909).

Diagnosis. Shell outline subtriangular to transversely elliptical. Biconvex, plicate shell with erect to sub-erect pedicle umbo, round foramen, and triangular-shaped delthyrium which may be partly covered with small deltidial plates. Pedicle valve with teeth supported by dental plates which terminate anteriorly at level of teeth. Brachial valve with median septum of variable length, height, and width, posteriorly thickens and forms base of notothyrial cavity in which elongate blade-like cardinal process is housed.

Morphological summary. Shell, commonly but not always, with width equal to or greater than length. Biconvex, plicate shell, commonly with brachial valve more convex than pedicle valve. Sulcus of variable form and size containing from 1 to 7 ribs. Each flank of pedicle valve occupied by 2 to 4 prominent ribs and 1 to 2 minor ribs. Pedicle umbo erect to suberect, foramen round and triangular delthyrium open or partly enclosed by small deltidial plates. Brachial valve contains median septum of variable length, height, and width (Table 1). Generally broadens posteriorly to form base of notothyrial cavity in which the elongate, blade-like cardinal process is housed. U-shaped sockets, which widen anterolaterally are bounded by elongate socket plates on inner edges. Triangular-shaped hinge plates curved ventrolaterally with crural bases at inner margins. Crus curve anterodorsally. In the pedicle valve the teeth are supported by dental plates which extend anteriorly to level of teeth before terminating sharply. Umbonal cavities are D-shaped.

Species assigned to Stegerhynchus. Since all of the species listed below have been described in detail elsewhere, only the pertinent remarks regarding each species is given. Representatives of the type material examined in detail are shown in Plate 18, and text-figs. 2 and 3. In the following descriptions the species are listed alphabetically.

Stegerhynchus angaciensis Chernyshev, 1937

Plate 18, figs. 46–48; text-fig. 2e, h

- 1937 *Stegerhynchus decemplicatus angaciensis* Chernyshev, p. 29; p. 71, pl. 1, figs. 15–18; text-figs. 1, 2.
 1955 *Stegerhynchus angaciensis* Borisyak, p. 46, pl. VI, figs. 9–11.
 1955 *Stegerhynchus angaciensis* var. *tryplicata* Borisyak, p. 47, pl. VI, figs. 6–8.
 1955 *Stegerhynchus angaciensis* var. *tryplicata* Borisyak; Zinchenko and Kulkov, 1960, p. 103, pl. S-28, fig. 7.
 1960 *Stegerhynchella angaciensis* (Chernyshev); Rzhonsnitskaya *et al.*, pl. 43, fig. 7a–d; text-figs. 246–247.
 1960 *Stegerhynchella angaciensis* (Chernyshev); Khalfina, p. 102, pl. S-28, fig. 5a–c.
 1970 *Stegerhynchus angaciensis* Chernyshev, 1937; Lenz, pp. 488–489, pl. 85, figs. 16–29.

Remarks. *S. decemplicatus angaciensis* Chernyshev, 1937 from the Late Silurian strata of western Mongolia and Tuva was used as the type species of the genus *Stegerhynchella* by Rzhonsnitskaya (1959). However, I follow Schmidt and McLaren (1965, p. H556) and Lenz (1970, p. 488) in considering that *Stegerhynchella* is probably a synonym of *Stegerhynchus*. Lenz (1970) assigned rhynchonellids from Prongs Creek to *S. angaciensis* (Pl. 18, figs. 46–48) even though these specimens have fewer ribs on their flanks than was considered diagnostic for *S. decemplicatus angaciensis*. This was done because inspection of Chernyshev's illustrations (1937, pl. 1, fig. 15a–d) showed specimens that are very similar to the Prongs Creek specimens. Comparison of the material used by Lenz (1970) with *S. decemplicatus angaciensis* from the River Elegest locality confirms this assessment. Like *S. praecursor*, *S. angaciensis* has a uniplicate sulcus. *S. angaciensis* differs from *S. praecursor* in having a higher brachial valve and a more elongate shell. These differences, however, may be ontogenetic or intraspecific in character.

Kulkov (1967, p. 76) listed *S. angaciensis* Borisyak, 1955 and *S. angaciensis* var. *triplicate* Borisyak, 1955 as synonyms of *F. borealis* (Schlothheim). These species would be best considered synonyms of *S. angaciensis* rather than *S. borealis*. *S. diodontia* can be separated from *borealis* by virtue of its distinctive uniplicate sulcus (Basset and Cocks, 1974, p. 26). It is extremely close to *S. angaciensis* Chernyshev, 1937; differing by being slightly more transverse and by having distinctive rugae.

Stegerhynchus? antiqua (Savage, 1913)

Plate 18, figs. 34–39

- 1913 *Camarotoechia? antiqua* Savage, p. 128, pl. 7, figs. 1 and 2.
 1964 *Ferganella? lincolnsis* Johnson, 1964; Johnson and Reso, p. 80, pl. 19, figs. 5–12.
 1970 *Ferganella* cf. *F. lincolnsis* Johnson, 1964; Lenz, pp. 487–488, pl. 85, figs. 1–15.
 1974 *Stegerhynchus? antiqua* (Savage, 1913); Amsden, p. 68, pl. 15, fig. 4a–i.
 1976 *Stegerhynchus* cf. *S. lincolnsis* (Johnson, 1964); Johnson *et al.*, pp. 64–65, pl. 47, figs. 1–12.

Remarks. Savage (1913) named *Camarotoechia? antiqua* for rhynchonellids from the Leemon Formation and the Noix Limestone of Illinois. Amsden (1974, p. 68) placed specimens UIX-872, 4784 with question in *Stegerhynchus* because the internal structures are unknown. Johnson and Reso (1964, p. 80) assigned rhynchonellids from the Sevy Dolomite of Nevada to the new species


EXPLANATION OF PLATE 18

- Figs. 1–15. *Stegerhynchus concinna* (Savage, 1913); 1–3, lateral, anterior, and pedicle views of the paratype, UI-RX-319; 4–6, lateral, anterior, and pedicle views of the paratype, UI-RX-316; 7–9, lateral, anterior, and pedicle views of the paratype, UI-RX-311; 10–12, lateral, anterior, and pedicle views of paratype, UI-RX-308; 13–15, lateral, anterior, and pedicle views of lectotype, UI-RX-855.
 Figs. 16–21. *Ferganella chattertoni* Lenz, 1970 (synonym of *C. concinna* herein); 16–18, lateral, anterior, and pedicle views of the holotype, GSC 48067; 19–21, lateral, anterior, and pedicle views of paratype, GSC 48068.
 Figs. 22–24. *Stegerhynchus? sp.* of Lenz (1970); lateral, anterior, and pedicle views, GSC 48066.
 Figs. 25–33. *Ferganella lincolnsis* Johnson and Reso, 1964; 25–27, lateral, anterior, and pedicle views of plaster cast of the holotype, UCLA 35050; 28–30, lateral, anterior, and pedicle views of plaster cast of UCLA 35054; 31–33, lateral, anterior, and pedicle views of a specimen from Prongs Creek assigned to *F. lincolnsis* Johnson and Reso by Lenz (1970).
 Figs. 34–39. *Stegerhynchus? antiqua* (Savage, 1913); 34–36, lateral, anterior, and pedicle views of paratype, UI-RX-872; 37–39, lateral, anterior, and pedicle views of paratype, UI-RX-4784.
 Figs. 40–45. *Stegerhynchus claritense* Amsden, 1968. 40–42, lateral, anterior, and pedicle views of paratype, OU 6385; 43–45, lateral, anterior, and pedicle views of holotype OU 6384.
 Figs. 46–48. *Stegerhynchus angaciensis* Chernyshev, 1937. Lateral, anterior, and pedicle views of a specimen from Prongs Creek assigned to this species by Lenz, 1970, GSC 25009.

All specimens $\times 2$.



JONES, brachiopod *Stegerhynchus*



TEXT-FIG. 3. Topotype specimens of *Stegerhynchus praecursor* from the type locality, collected by Foerste, Clinton Bed, Clifton, Tennessee. Figs. a-e and f-j, pedicle, brachial, lateral, anterior, and posterior views of USNM 218094 and USNM 218098 respectively. Note cardinal process evident in figs. d and i. Fig. k, brachial valve of USNM 304184 showing extent of median elevation. Fig. l, brachial valve of USNM 218092 showing median elevation that is longer and more strongly developed than that in USNM 304184. Figs. m and n, pedicle valve of USNM 304185 and USNM 304186 showing dental lamellae.

lincolnensis (Pl. 18, figs. 25-30). Although the internal structures of these specimens had been obliterated by silification Johnson tentatively assigned the species to *Ferganella*. However, Johnson *et al.* (1976, p. 64) later transferred *lincolnensis* to *Stegerhynchus*, claiming that the internal structures of specimens from the Roberts Mountains Formation of central Nevada were more indicative of *Stegerhynchus* than *Ferganella* as defined by those authors.

Comparison of the holotype of *lincolnensis* (UCLA 35050, Pl. 18, figs. 25-27) with the lectotype of *antiqua* (UI-RX-872, Pl. 18, figs. 34-36) shows that they have similar shell outline, similar profile, and similar ribbing both in the sulcus and on the flanks of the shell. More important, both have a sulcus that is poorly defined and developed only in the anterior portion of the shell. The internal structures of both *lincolnensis* and *antiqua* remain unknown. In spite of their morphological similarity *antiqua* has been reported only from upper Ordovician strata whereas *lincolnensis* has been reported only from Ludlovian strata (text-fig. 4).

Pending discovery of internal structures *lincolnensis* is provisionally placed as a synonym of *antiqua*. *S.?* *antiqua* is separated from *S. concinna* because it has a poorly defined sulcus which is well defined in the latter. *S.?* *antiqua* has an unequally convex shell and this serves to distinguish it from other species of *Stegerhynchus* which have a brachial valve that is more convex than the pedicle valve.

Stegerhynchus borealis (Von Buch, 1834)

- 1822 *Anomia Terebratula lacunosus* Linnaeus; Schlotheim, pl. 20, fig. 6a-c, non Linnaeus, 1758.
 1832 *Terebratula borealis* Schlotheim, p. 65 nomen nudum.
 1834 *Terebratula borealis* Von Buch.
 1869 *Rhynchonella borealis* (Schlotheim); Davidson, p. 174, pl. 21, figs. 14, 15, 17, 19, 20, and 24-27.
 1937 *Camarotoechia borealis* (Von Buch); St. Joseph, p. 33, figs. 1-5.
 1937 *Ferganella* cf. *F. borealis* (Schlotheim); Nikiforova, p. 42, pl. 6, fig. 17a-d.
 1954 *Camarotoechia(?) borealis* (Schlotheim); Nikiforova, p. 98, pl. X, fig. 4.

- 1967 *Ferganella borealis* (Schlotheim); Kulkov, pp. 76-79, fig. 31; pl. XII, figs. 7-11. *non Ferganella? lincolnensis* Johnson and Reso, 1964, p. 80, pl. 19, figs. 5-12.
- 1974 *Ferganella borealis* (Von Buch); Bassett and Cocks, p. 26.
- 1976 *Ferganella borealis* (Schlotheim); Sheehan, p. 729, pl. 5, figs. 1-5.
- 1976 *Ferganella* cf. *F. turkestanica* Nikiforova, 1937; Smith, p. 29, text-fig. 19; pl. 7, figs. 20-25.
- 1977 *Ferganella chattertoni* Lenz, p. 1542, pl. 8, figs. 1-16.

Remarks. In the literature there is some confusion over the authorship of the species *borealis*. Kulkov (1967, p. 76) and Sheehan (1976, p. 729) both attributed *borealis* to Schlotheim (1832) whereas Bassett and Cocks (1974, p. 26) attributed it to Von Buch, 1834. Schlotheim (1832, p. 65) used the name *Terebratula borealis* without illustrating or describing the specimens he was assigning to the species. Thus, the name remained a *nomen nudum* until Von Buch (1834) provided an adequate description of *borealis*. The species should therefore be assigned to Von Buch, 1834 as suggested by Bassett and Cocks (1974, p. 26). Although St. Joseph (1937, pp. 33 and 45) considered that Von Buch (1834) had formerly chosen the type specimen of the species, Bassett and Cocks (1974, p. 26) have shown that this is not the case. Thus, the description by St. Joseph (1937, p. 33) of the specimen figured by Schlotheim (1822, pl. 20, fig. 6a-c) should be regarded as the first description of the lectotype of the species (Bassett and Cocks, 1974, p. 26). The available descriptions suggest that *S. borealis* has a high degree of intraspecific variation. The problem is one of defining the exact range of variation for the species. For example, St. Joseph (1937, p. 46) considered *diodonta* Dalman, 1828 as a variety of *borealis* whereas Bassett and Cocks (1974, p. 26) considered them two separate species. Similarly, Kulkov (1967, p. 76) considered *F.? lincolnensis* Johnson, 1964 a synonym of *borealis* while Sheehan (1976, p. 729) maintained the two as separate species. Lenz (1977, p. 1452) compared *F. chattertoni* with *F. borealis*, which Sheehan (1976) described from Utah. He considered the two distinct because the Mackenzie Mountains material lacked a deep, ventral muscle field, had 'normal' sized costae on either side of the sulcus and possibly a sharper, more pointed beak. Comparison of the type material of *chattertoni* with Sheehan's illustrations of *F. borealis* shows that these differences are probably valid. However, comparison of *chattertoni* with *borealis* as described by Kulkov (1967) and St. Joseph (1937) strongly suggests that *chattertoni* falls within the range of morphological variation of *borealis*. For this reason *chattertoni* is included as a synonym of *borealis*.

Stegerhynchus claritense Amsden, 1968

Plate 18, figs. 40-45

- 1968 *Stegerhynchus claritense* Amsden, pp. 61-63, text-fig. 47, Table 32; pl. 17, fig. 1a-k.

Remarks. *S. claritense* was named by Amsden (1968, pp. 61-62) for specimens from the Fitzhugh Member of the Clarita Formation of Arkansas. It is separated from other species of *Stegerhynchus* by virtue of its very distinctive subrounded ribs as compared to the angular ribs of other species.

Stegerhynchus concinna (Savage, 1913)

Plate 18, figs. 1-15; text-fig. 2a, b, c

- 1913 *Camarotoechia? concinna* Savage, pp. 128-129, pl. 7, fig. 3 (*non* Billings, 1866).
- 1974 *Stegerhynchus concinna* (Savage); Amsden, pp. 66-68, text-figs. 41-42, Table 9; pl. 14, figs. 3, 4; pl. 15, figs. 1-3.

Remarks. Since the original specimens of *concinna* Savage (1913) from the Edgewood Formation in Pike County, Missouri, and near Thebes, Alexander County, Illinois, could not be found, Amsden (1974, p. 67) designated specimen UI-RX-855 as the lectotype of the species. *S. concinna* differs from *S. praecursor* because it has more ribs (1 to 3 compared to 1) in its sulcus. It has subequally convex valves, whereas *S. praecursor* has a brachial valve that is more convex than the pedicle valve.

S. concinna was separated by Lenz (1977, p. 1425) from *F. chattertoni* because of its more distinctly rounded outline, fewer fold and sulcus costae, and its long median septum. However, comparison of the type series of the two species (Pl. 1, figs. 1–15 and 16–21) shows that *chattertoni* has three ribs in the sulcus while *concinna* has one to three ribs in the sulcus (Table 1). Thus, separation on this basis is not always possible. Although *F. concinna* is generally rounder than *chattertoni*, some of the more extreme forms have a very similar outline to *chattertoni*. The median septum in *chattertoni* is 36 to 40% of valve length (based on three specimens) while in *concinna* the septum is 55 to 65% of valve length (based on two specimens). However, this apparent difference in the length of the median septum should be treated with caution since study of other species of *Stegerhynchus* has shown that there can be a wide range of variation in the length of this feature even amongst shells from the same locality.

Stegerhynchus praecursor Foerste, 1909

Text-fig. 3

- 1909 *Stegerhynchus whitii-praecursor* Foerste, pp. 96–98, pl. III, figs. 47A, B, C.
 1944 *Stegerhynchus whitii-praecursor* Foerste, 1909; Shimer and Shrock, p. 309.
 1955 *Ferganella praecursor* (Foerste); Cooper, p. 55.
 1965 *Stegerhynchus praecursor* Foerste, 1909; Schmidt and McLaren, p. H556.
 1978 *Stegerhynchus cliftonensis* Foerste, 1909; Amsden, p. 28, pl. 11, figs. 1–9.

Remarks. The specimens of *S. cliftonensis* and *S. praecursor* figured by Foerste (1909, pl. III, figs. 47 and 48) are not available for further study. According to the original descriptions of the species (Foerste, 1909, pp. 96–98) they were considered to differ only in that *S. cliftonensis* had three ribs in its sulcus while *S. praecursor* had one rib in its sulcus. Both species were based on material from the Clinton Bed at Clifton, Tennessee (Foerste, 1909, p. 97). In the USNM collection there is a topotypic set of thirty specimens (collected by Foerste) which are labelled as *S. praecursor*. However, as noted by Amsden (1978, p. 23), most of these shells have three ribs in the sulcus and are thus more allied to *S. cliftonensis* as originally described by Foerste (1909, p. 97). Although Amsden (1978, p. 23) assigned these shells to *S. cliftonensis*, he noted that they could well be variants of *S. praecursor* rather than a separate species. The latter possibility seems more feasible since some shells in the topotypic collection have two ribs in the sulcus and could therefore be assigned to neither *S. cliftonensis* nor *S. praecursor* if Foerste's original definitions were followed. Rather, there is the implication that this particular attribute alone is not very useful for separating species. This point is emphasized by the fact that other species, such as *S. borealis*, have anywhere from one to seven ribs in the sulcus. In view of these points *S. cliftonensis* is herein included as a synonym of *S. praecursor*. The available specimens from the Clinton Bed of Tennessee clearly show that *Stegerhynchus* has a well-defined median septum which tends for 50 to 60% of the length of the dorsal valve. The notothelial cavity houses a long, narrow cardinal process (text-fig. 3).

EXPLANATION OF PLATE 19

Figs. 1–18, 25–33, 40–42. *Stegerhynchus borealis* (von Buch, 1834); lateral, anterior, and pedicle views of a series of specimens from assemblage M113 collected at Cape Admiral M'Clintock, Read Bay Formation, Somerset Island; UA3701 to UA3709 inclusive.

Figs. 19–24, 34–39, 43–45. *Stegerhynchus angaciensis* Chernyshev, 1937; lateral, anterior, and pedicle views of specimens from basal unit of Member C of the Read Bay Formation at Goodsir Creek, Cornwallis Island, UA3710 to UA3714 inclusive.




JONES, brachiopod *Stegerhynchus*

CONSTITUTION AND AGE OF *STEGERHYNCHUS**Species wrongly assigned to Stegerhynchus*


In addition to the above species assigned to *Stegerhynchus* there are a number of species that have been wrongly assigned to the genus. In almost every case this resulted from confusion over the true type species of *Stegerhynchus*. Thus, Havlicek (1961, pp. 80-91) assigned *daphne* Barrande, 1847, *infelix* Barrande, 1879, *incohans* Havlicek, 1961, *nympha* Barrande, 1847, and *pseudolivonicus* Barrande, 1847 to *Stegerhynchus* while Kulkov (1963, pp. 45-50) assigned *daphne* Barrande, 1879, *nympha* Barrande, 1879, and possibly *dichotoma* Khalfin, 1948 to *Stegerhynchus*. In his discussion of the genus, Kulkov (1963, pp. 45-50) also assigned *pseudolivonica* Barrande to *Stegerhynchus*. These assignments by Havlicek (1961) and Kulkov (1963) were made because they believed that *R. whitii* Hall was the type species of the genus as argued by Cooper (1955, pp. 54-55). None of these species possess a cardinal process and therefore cannot belong to *Stegerhynchus*. The affinity of these species is uncertain and requires further study before a definite assignment can be made. In 1967 Kulkov assigned *nuculus* Sowerby, 1839 to *Stegerhynchus*. However, inspection of the pertinent figures shows that the specimens sectioned do not contain a cardinal process (Kulkov, 1967, p. 83) and cannot, therefore be assigned to *Stegerhynchus*. Shimer and Shrock (1944, p. 309) also listed the species *whitii*, Hall, *neglectum*, Hall, *indianense*, Hall, and *acinus*, Hall as members of *Stegerhynchus*. However, no information about internal structures was supplied; therefore, it is impossible to verify the validity of this assessment.

Age ranges of Stegerhynchus

Apart from isolated occurrences in the uppermost Ordovician of Missouri, *Stegerhynchus* is restricted to Silurian strata (text-fig. 4). Species such as *S. borealis* and *S. diodonta* have relatively long ranges while other species such as *S. diodonta* are relatively restricted. Apparently, *Stegerhynchus* does not occur in Devonian strata. *S. borealis* occurs in the Read Bay Formation of Arctic Canada in strata of upper Ludlovian age and this extends the range of the species as known from the literature. In the basal part of Member C of the Read Bay Formation at Goodsir Creek (text-fig. 5) *S. borealis* and *S. angaciensis* occur together thereby confirming the overlapping age ranges of these species.



TEXT-FIG. 4. Geographic and stratigraphic distribution of species assigned to *Stegerhynchus*.



TEXT-FIG. 5. Geographic and stratigraphic locations of *Stegerhynchus borealis* and *S. angaciensis* in the Read Bay Formation of Somerset, Prince of Wales, and Cornwallis Islands. *S. borealis* is relatively common, especially in the lower part of the formation whereas *S. angaciensis* is relatively rare, occurring only in the basal part of Member C on eastern Cornwallis Island.

RHYNCHONELLIDS FROM THE READ BAY FORMATION
OF ARCTIC CANADA

The Upper Silurian Read Bay Formation of Somerset, Prince of Wales and Cornwallis Islands of Arctic Canada, contains an abundant fauna of brachiopods. Rhynchonellids occur at many levels in the Read Bay Formation of these islands. *S. borealis* was collected from six localities on Somerset Island, three localities on Prince of Wales Island, and from members A and C of the type section of the Read Bay Formation at Goodsir Creek on Cornwallis Island. *S. angaciensis* has only been recorded from four localities on the east coast of Cornwallis Island (text-fig. 5).

Stegerhynchus borealis (Von Buch, 1834)

Plate 19, figs. 1-18, 22-30, 37-39; Plate 20


Horizon and age. *S. borealis* occurs at many horizons in the Read Bay Formation but is commonest in the lower part of the formation (text-fig. 5). Thus, it would appear to have a Ludlovian to Pridolian range on Somerset, Prince of Wales, and Cornwallis Islands.

Preservation. Most *S. borealis* in the Read Bay Formation are preserved as complete calcareous shells. However, silicified specimens have been recovered from localities on Prince of Wales and Somerset Islands and the description of the internal structures is based on this material.


Description. External morphology. *S. borealis* has a biconvex, plicate shell that attains a maximum length of about 2 cm. In small specimens (<1 cm long) the valves are subequally convex (Pl. 19, figs. 37-39) whereas in larger, more mature shells the brachial valve is more convex than the pedicle valve (Pl. 19, figs. 1-15). Pedicle umbo generally erect. Delthyrium triangular and commonly bordered by small deltidial plates. Shell outline subtriangular with shell width generally exceeding shell length (text-fig. 7a). Sulcus commences 4 to 5 mm from the pedicle beak and becomes wider and deeper anteriorly (Pl. 19, figs. 2, 5, and 8). Most commonly, the sulcus contains three angular ribs (Pl. 19, figs. 5, 14, 23, and 26) but there is a wide range of variation about this mode (Pl. 19, figs. 2, 8, 11, and 29). For example, 80% of the shells in assemblage M113 have 3 ribs in the sulcus while the remaining 20% have 1, 2, 4, 6, or 7 ribs (text-fig. 6a). The number of ribs in the sulcus does not appear to be a function of ontogeny since there is no relationship between shell size and the number of ribs in the sulcus (text-fig. 7c). Each shell flank has 4 to 8 ribs which fade posteriorly (Pl. 19, fig. 3). Apical angle ranges from 66 to 95 degrees (text-fig. 6c).

On the pedicle valve slightly elongate and relatively stout teeth (Pl. 20, fig. 8) are supported by dental lamellae that diverge slightly anterolaterally (Pl. 20, fig. 4). The dental plates, which terminate abruptly, are short, being confined to the umbonal region of the shell (Pl. 20, fig. 4). Umbonal cavities D-shaped. On the brachial valve the sockets widen anterolaterally (Pl. 20, fig. 2) and are bounded by blade-like dental socket plates (Pl. 20, figs. 2 and 6). The triangular-shaped hinge plates are convex anterolaterally and have crural bases on their inner edges. Crus extend anteroverally (Pl. 20, fig. 3). The notothyrial cavity is relatively large and has as its base the notothyrial platform which supports an elongate, blade-like cardinal process (Pl. 20, figs. 1, 2, and 7). The notothyrial platform is supported by a median septum which becomes narrower and lower anteriorly, eventually disappearing at about midlength (Pl. 20, fig. 6).

Remarks. These Read Bay rhynchonellids are assigned to *S. borealis* because they are close to *S. borealis* as described by Bassett and Cocks (1974, p. 26), St. Joseph (1937), Kulkov (1967, pp. 76-79), and Sheehan (1976, p. 728). They are also close to specimens Br 103764 and Br 103765 of *S. borealis*. Smith (1976) assigned rhynchonellids from the Upper Silurian Douro Formation (equivalent to the Read Bay Formation of Somerset Island and Member A of the Read Bay Formation on Cornwallis Island) to *F. turkestanica*. However, these specimens have internal structures more typical of *Stegerhynchus*. They are very similar to the specimens described as *S. borealis* in this paper.



TEXT-FIG. 6. Comparison of (a) number of ribs in sulcus, (b) length, (c) width, (d) height, (e) apical angle, and (f) number of ribs on each shell flank of pedicle valve for assemblage M113 of *Stegerhynchus borealis* and *S. angaciensis* from the east coast of Cornwallis Island. The parameters are defined in fig. g. Note that the major difference between the two species is in the number of ribs in the sulcus.



TEXT-FIG. 7. Bivariate graphs comparing *Stegerhynchus angaciensis* (from east coast of Cornwallis Island) and *S. borealis* (assemblage M113) for (a) length versus width and (b) width versus height demonstrating that the two species are very similar in these respects. (c) pictogram showing the absence of any relationship between the number of ribs in the sulcus and shell size of *S. borealis*.

Stegerhynchus angaciensis Chernyshev, 1937

Plate 19, figs. 16–21, 31–36, and 40–42

Occurrence and age. *S. angaciensis* has only been found in restricted numbers at four localities on eastern Cornwallis Island. *S. angaciensis* occurs in the basal unit of Member C of the Read Bay Formation at Goodsir Creek (text-fig. 5). At localities (2, 3, and 4) to the south and north of Goodsir Creek, *S. angaciensis* has only been found in the scree. However, their position in the scree and the associated fauna clearly points to them originating in the same stratigraphic position as the specimens at Goodsir Creek. Their occurrence at the base of Member C strongly suggests a Late Ludlovian age for the brachiopods since conodonts from Members A and C suggest that Members A, B, and part of Member C are of Ludlovian age while the upper part of Member C is of Prividian age (Uyeno, 1977, fig. 41.2).

Preservation. All of the specimens of *S. angaciensis* from eastern Cornwallis Island are preserved as articulated calcareous shells. Most of the shells are complete.

Description. Biconvex, plicate shell that attains a maximum length of about 2 cm. Brachial valve more convex than pedicle valve even in small specimens, and this becomes even more pronounced in larger, more mature specimens. Pedicle umbo pointed, erect to sub-erect. Pedicle foramen is clearly visible. Shell outline subtriangular. In small specimens (<1 cm) width is approximately equal to length; however, in larger, more mature specimens width generally exceeds length. Well-defined uniplicate sulcus begins 2 to 3 mm from beak and is laterally bounded by steep-sided sulcus walls (Pl. 19, figs. 17, 20, 32, and 35). The brachial valve has a correspondingly strong fold with two angular ribs. Shell flanks are occupied by 5 to 6 strong, angular ribs and 1 to 2 minor angular ribs.

Internal structures resemble those of *S. borealis*.

Remarks. The specimens of *S. angaciensis* from Cornwallis Island are very similar to those described and illustrated by Chernyshev (1937, pl. 1).

Acknowledgements. I acknowledge the support of grant A-5121 to Dr. O. A. Dixon from the National Research Council of Canada; field-work at Goodsir Creek in 1977 was supported by the Boreal Institute, University of Alberta. The Polar Continental Shelf Project provided logistic aid for all of the field-work. I am indebted to Dr. T. E. Bolton, Geological Survey of Canada; Dr. T. W. Amsden, Oklahoma Geological Survey; Dr. T. Blake, University of Illinois and Ms. Louella Saul, University of California, Los Angeles for making type material available for this study; S. R. Williams provided additional material from Prince of Wales Island. I thank F. Dimitrov for drafting, G. Braybrook for SEM photography, C. van der Lee for assistance in preparatory work, Drs. C. Stelck, B. D. E. Chatterton, M. G. Bassett, and T. W. Amsden for additional help.

EXPLANATION OF PLATE 20


Figs. 1–8. *Stegerhynchus borealis* (von Buch, 1834); 1, Posterior portion of articulated shell showing the teeth in sockets, triangular-shaped hinge plates, and relatively large notothyrial cavity housing elongate, blade-like cardinal process. $\times 22$. 2, Interior of brachial valve showing curved hinge plates, notothyrial cavity, elongate cardinal process, median septum becoming lower and narrower anteriorly, and muscle scars on either side of the septum. Note teeth broken off into sockets. $\times 20$. 3, Posterior portion of brachial valve showing crus extending from hinge plates. $\times 23$. 4, View of posterior portion of pedicle valve from anterior showing dental plates and D-shaped umbonal cavities. This valve matches the brachial valve shown in fig. 2. $\times 19$. 5, Posterior portion of articulated shell. $\times 29$. 6, Interior of brachial valve showing full extent of median septum. Note that septum is well developed even in this small shell. $\times 10$. 7, View of the notothyrial cavity from anterior margin showing shape of cavity, shape of cardinal process, and height of cardinal process above cavity floor. $\times 26$. 8, Enlarged view of tooth from pedicle valve. $\times 47$.




1



2



3



4



5



6



7



8

REFERENCES

- AGER, D. V. 1965. Mesozoic and Cenozoic rhynchonellacea. In: *Treatise on Invertebrate Paleontology*, Pt. H, Brachiopoda: H597-632, University of Kansas Press.
- ALEKSEVA, R. E., GRATSIANOVA, R. T., ELKIN, E. A. and KULKOV, N. P. 1970. Stratigrafiya i brachiopoda nizhnego Devona severo vostochnogo Salaira. *Tr. Inst. Geol. Geofiz. Sibirsk. Otd. Akad. Nauk. S.S.S.R.* **72**, 1-188.
- AMSDEN, T. W. 1968. Articulate brachiopods of the St. Clair Limestone (Silurian), Arkansas, and the Clarita Formation (Silurian), Oklahoma. *J. Paleont.* (Mem. 1), **42**, 1-99.
- 1974. Late Ordovician and Early Silurian articulate brachiopods from Oklahoma, southwestern Illinois, and eastern Missouri. *Bull. Okla. Geol. Surv.* **119**, 84 pp.
- 1978. Articulate brachiopods of the Quarry Mountain Formation (Silurian), eastern Oklahoma. *Ibid.* **125**, 75 pp.
- and BOUCOT, A. J. 1958. Stratigraphy and Paleontology of the Hinton Group in the Arbuckle Mountain Region. *Ibid.* **78**, 199 pp.
- BARRANDE, J. 1847. *Über die Brachiopoden der Silurischen Schichten von Boehmen*. 104 pp. Vienna.
- 1879. *Système Silurien du centre de la Bohême*. 1^{re} partie, vol. 5, Classe des Mollusques. Orde des Brachiopodes. 226 pp. Prague, Paris.
- BASSETT, M. G. and COCKS, L. R. M. 1974. A review of Silurian Brachiopods from Gotland. *Fossils Strata*, **3**, 56 pp.
- BILLINGS, E. 1866. *Catalogues of the Silurian fossils of the island of Anticosti with descriptions of some new genera and species*. Canada Geol. Surv. Spec. Publ. 93 pp. Ottawa.
- BORISYAK, M. A. 1955. Siluriiskikh (Wenlockski) Brachiopody i Karagandinskoy oblasti. (Silurian (Wenlock) brachiopods of the Karagandinskoy region.) *Materialy VSEGEI*, Novaya Seria, **5**, no. 2.
- CHERNYSHEV, T. N. 1937. Siluriiskie brachiopody Mongolii i Tuvy. *Akad. Nauk USSR, Trudy Mongolskoi Kouissii*, **29**, 1-94.
- COCKS, L. R. M. 1978. A review of British Lower Paleozoic brachiopods, including a synoptic revision of Davidson's monograph. *Palaentogr. Soc. (Monogr.)*, 256 pp.
- COOPER, G. A. 1955. New Genera of middle Paleozoic brachiopods. *J. Paleont.* **29**, 45-63.
- DAVIDSON, T. 1869. A monograph of the British fossil Brachiopoda. Part 7. Silurian. *Palaentogr. Soc. [Monogr.]*, pp. 169-248.
- DROT, J. 1964. Rhynchonelloidea et Spiriferoidea Siluro-Devonien du Maroc pre-Saharien. *Notes Mém. Serv. geol. Maroc*, **178**, 1-287.
- FOERSTE, A. F. 1909. Fossils from the Silurian Formations of Tennessee, Indiana and Kentucky. *Bull. Denison Univ. Sci. Lab.* **14**, 61-116.
- GRATSIANOVA, R. T. 1967. Brachiopody i stratigrafiya nizhnego Devona Gornogo Altaiya. *Akad. Nauk. SSSR, Sibirsk. Otdel., Inst. Geol. i Geofiz.*, pp. 1-160.
- HALL, J. and CLARKE, J. M. 1894. An introduction to the study of Brachiopoda. *13th Ann. Rept. N.Y. St. Geol. for 1893*. Paleont., pt. 2, 751-943.
- HAVLICEK, J. P. 1961. Rhynchonelloidea des bohmischen alteren Palaozoikum (Brachiopoda). *Rozpr. ustred. Ust. Geol.* **27**, 1-211.
- JOHNSON, J. G. and RESO, A. 1964. Probable Ludlovian brachiopods from the Sevy Dolomite of Nevada. *J. Paleont.* **38**, 74-84.
- BOUCOT, A. J. and MURPHY, M. A. 1976. Wenlockian and Ludlovian Age brachiopods from the Roberts Mountains Formation of central Nevada. *Univ. Cal. Publ. Geol. Sci.* **115**, 102 pp.
- KHALFIN, L. L. 1948. Fauna i Stratigrafiya devonskikh otlozhenii Gornovo Altai. (Fauna and stratigraphy of Devonian deposits of Altai.) *Izv. Tomsk. Polytechni. In-Ta.* **65**, vyp 1.
- KHALFINA, V. K. 1960. Biostratigrafiya paleozoya Sayano-Altaijskogo gornogi oblasti. *Trudy Sibirsk Nauchuo-Issled. Inst.* **20**, 99-109.
- KOZLOWSKI, ROMAN. 1929. Les Brachiopodes Gothlandiens de la Podolie Polonaise. *Paleont. Polon.* **1**, 254 pp.
- KULKOV, N. P. 1963. *Brachiopody Solovikhinskikh sloev nizhnego Devona gornogo Altaya*. Akad. Nauk USSR, Moscow. 131 pp.
- 1967. *Brachiopody i Stratigrafiya Silura gornogo Altaya*. Akad. Nauk. USSR, Moscow. 151 pp.
- LENZ, A. C. 1970. Late Silurian brachiopods of Prongs Creek, northern Yukon. *J. Paleont.* **44**, 480-500.
- 1977. Llandoveryan and Wenlockian brachiopods from the Canadian Cordillera. *Cau. J. Earth Sci.* **14**, 1521-1554.
- LINNAEUS, C. 1758-1767. *Systema Naturae sive Regna tria Naturae systematicae proposita per Classes, Ordines, Genera et Species*. 10th edn., 1758, 1-284. Uppsala.

- NIKIFOROVA, O. I. 1937. Upper Silurian Brachiopoda of the central Asiatic part of U.S.S.R. *Paleontology of U.S.S.R. Monogr.* **35** (1), 94 pp.
- RZHONSNITAKAYA, M. A., LIKHAREV, B. K. and MAKRIDIN, V. P. 1960. Otryad Rhynchonellida. In T. G. SARYCHEVA (ed.), *Osnovy Paleontologii Mshanki, Brakhiopody*, pp. 239-257.
- 1959. K sistematike rinkhonellid. *Paleont. Zh.* **1**, 25-35.
- SAVAGE, T. E. 1913. Stratigraphy and paleontology of the Alexandrian Series in Illinois and Missouri. *Bull. Ill. St. Geol. Surv.* **23**, 67-160.
- SCHLOTHEIM, E. F. 1822. *Nachtrage zur Petrefactenkinde*, I-XI. 100 pp., atlas 21. Gotha.
- 1832. *Systematisches Verzeichniss der Pet Sammling*, **65**, 88 pp. Gothenburg.
- SCHMIDT, H. and McLAREN, D. J. 1965. Paleozoic Rhynchonellacea. In: *Treatise on Invertebrate Paleontology*, Pt. H. Brachiopoda, pp. H552-597. University of Kansas Press.
- SCHUCHERT, C. and LEVENE, C. M. 1929. Brachiopoda (Generum et genotyporum index et bibliographica), *Fossilium Catalogus*, **1**, Animalia, 140 pp. Junk (Berlin).
- SHEEHAN, P. M. 1976. Llandoveryan and Wenlockian brachiopods from the Canadian Cordillera. *Can. J. Earth Sci.* **14**, 1521-1554.
- SHIMER, H. W. and SHROCK, R. R. 1944. *Index fossils of North America*. Mass. Inst. Tech. (Cambridge, Mass.). 837 pp.
- SMITH, R. E. 1976. Biostratigraphy and paleoecology of the *Atrypella* Community, Upper Silurian Douro Formation, Devon Island, District of Franklin. In Contributions to Canadian Paleontology. *Bull. Geol. Surv. Can.* **256**, 13-31.
- ST. JOSEPH, J. K. S. 1937. On *Camarotoechia borealis* (Von Buch, 1834, ex. Schlotheim, 1832). *Geol. Mag.* **74**, 33-48.
- UYENO, T. T. 1977. Summary of conodont biostratigraphy of the Read Bay Formation at its type section and adjacent areas, eastern Cornwallis Island, District of Franklin. In Report of Activities, Part B, *Pap. Geol. Surv. Can.* **77-1B**, 211-216.
- VON BUCH, H. 1834. Über Terebrateln. *Abh. Dt. Akad. Wiss. Berl.* (for 1883), pp. 21-144.
- ZINCHENKO, V. G. and KULKOV, J. P. 1960. Plechenogye silurii v kn: Biostratografiya Sayano Atlaskoi gornoy oblasti. (Silurian Brachiopods: in Paleozoic Stratigraphy of the upper Altai region.) *Trudy SNIIGGIMS*, **20**.

B. JONES

Paleontological Collections
Department of Geology
University of Alberta
Edmonton
Canada T6G 2E3

Typescript received 14 August 1979

Revised typescript received 20 February 1980